



Egyptian Herbal Monograph

Egyptian Herbal Monograph

Volume 3

Herbal Formulations Used in Egypt

Egyptian Drug Authority (EDA)

2026





Egyptian Herbal Monograph

Medicinal Plants Used in Egypt

Anise oil, Fennel oil, Licorice, Thyme زيت الينسون / زيت الشمر / عرقسوس / زعتر

1. Names & Synonyms

Anise (1)

***Pimpinella anisum* L.**

Family: Apiaceae (Umbelliferae)

Syns: *Anisum officinarum* Moench., *A. vulgare* Gaertn., *Apium anisum* L. Crantz., *Carum anisum* L. Baill., *Pimpinella anisum* cultum Alef., *P. aromatica* Bieb., *Selinum anisum* L. E.H.L. Krause., *Sison anisum* Spreng., *Tragium anisum* Link.

Arabic: Yansoon ينسون

English: Anise, Aniseed.

Fennel (2)

***Foeniculum vulgare* Mill.**

Family: Apiaceae (Umbelliferae).

Syns: *Anethum foeniculum* Clairv., *A. foeniculum* L., *A. rupestre* Salisb., *Feniculum commune* Bubani., *F. azoricum* Mill., *F. capillaceum* Gilib., *F. dulce* DC., *F. foeniculum* L. H. Karst., *F. officinale* All., *F. panmorium* DC., *F. piperitum* DC., *F. sativum* Bertol., *Ligusticum divaricatum* Hoffmannsegg et Link, L., *Foeniculum* Crantz., *Meum foeniculum* L. Spreng., *Ozodia foeniculacea* Wight et Arn., *Selinum foeniculum* L. E.H.L.Krause.

Arabic: Shamar شمر

English: Fennel

Licorice (3)

***Glycyrrhiza glabra* L. and its varieties or *Glycyrrhiza uralensis* Fisch.**

Syn. *Liquiritiae officinalis* Moench.



Family: Fabaceae (Leguminosae).

Arabic: Erqsos, Erqsos عرقسوس

English: licorice, licorice root, liquorice, liquorice root, sweet root and sweet wood.

Thyme

***Thymus vulgaris* L. (4)**

Syns.: *Origanum thymus* Kuntze, *Thymus collinus* Salisb.

Family: Lamiaceae (Labiatae).

Arabic: Za'ater زعتر

English: English Thyme, Garden Thyme.

***Thymus zygis* L. (5)**

Syns.: *Origanum zygis* (L.) Kuntze, *Thymus angustifolius* Salisb.

Family: Lamiaceae (Labiatae).

Arabic: Za'ater زعتر

English: Spanish Thyme.

2. Parts used for medicinal purpose

Anise: Dried ripe fruits (1).

Fennel: Dried ripe fruits (6).

Licorice: Dried Root (7-10) and rhizome (7, 10).

Thyme: Dried and fresh herb (11,12).

3. Major chemical constituents

Anise oil (1)

contains mainly trans-anethole, estragole (methylchavicol, isoanethole), β -anisaldehyde, α -terpineol, cis-anethole.

Fennel oil (13)

trans-anethole (+)-fenchone, estragole (methylchavicol), limonene, p-anisaldehyde, α -pinene and α -phellandrene.



Licorice (14)

- **Saponins:** Glycyrrhizin and glycyrrhizic acid (glycyrrhizinic acid), as a mixture of potassium and calcium salts.
- **Flavonoids:** glycosides of liquiritigenin and isoliquiritigenin, such as liquiritin, isoliquiritin, liquiritin apioside and licuraside (15).
- **Others:** Essential oil (mainly geraniol and geranyl hexanolate) and sterols (β -sitosterol, dihydrostigmasterol) (15).

Thyme

- **Essential oil :**the main components are thymol, carvacrol, *p*-cymene, α and β -terpinene, linalool, terpinen-4-ol, borneol, 1,8- cineole, α -thujene, α -pinene, and caryophyllene (11).
- **Flavonoids:** apigenin, narigenin, kaempferol, and luteolin (and its glycosides) (16).
- **Phenolic acids:** salvianolic, rosmarinic, cinnamic, ferulic, caffeic and gallic acids (16).
- **Others:** monoterpene glycosides.

4. Medicinal Uses (Indications)

- Expectorant in cough associated with cold (11,6,7,8,10,17,18).
- Relief of upper respiratory tract catarrh (17, 19-21).

5. Herbal preparations correlated to medicinal use

Combination of anise oil, fennel oil, licorice powder extract (extraction solvent: water, alcohol or hydro-alcoholic), and thyme aqueous extract.

Herbal preparations are in pharmaceutical dosage forms. The pharmaceutical form should be described by the pharmacopoeia full standard term.

6. Posology and method of administration correlated to medicinal use



Adults: Combination of 7.5 mg of anise oil, 0.15 mg of fennel oil, 1200 mg of licorice extract, and 750 mg of thyme extract, 3 times daily.

Adolescents (12–18 years): Combination of 2.5 mg of anise oil, 0.05 mg of fennel oil, 400 mg of licorice extract, and 250 mg of thyme extract, 3 times daily.

Children (8–12 years): Combination of 2.5 mg of anise oil, 0.05 mg of fennel oil, 400 mg of licorice extract, and 250 mg of thyme extract, twice daily.

Children (4–8 years): Combination of 1.25 mg of anise oil, 0.025 mg of fennel oil, 200 mg of licorice extract, and 125 mg of thyme extract, twice daily.

Duration of use: Not to be used more than one week. If the symptoms persist longer than one week, a doctor or a pharmacist should be consulted.

Method of administration: Oral use.

7. Contraindications.

- Hypersensitivity to the active substances, to any other plants of the same families.
- Cholestatic liver disorders, liver cirrhosis, hypertonia, hypokalemia, severe kidney insufficiency and cardiovascular-related disorder, because licorice ingestion has resulted in symptoms of primary hyperaldosteronism, such as water and sodium retention and hypokalemia (22).
- Not to be used concomitantly with diuretics, cardiac glycosides, corticosteroids, stimulant laxatives or other medications which may aggravate electrolyte imbalance (7,10,23).

8. Special warnings and precautions for use

- If the symptoms worsen or if dyspnoea, fever or purulent sputum occurs during the use of the medicinal product, a doctor or a pharmacist should be consulted (11,16,19).



- This medication is not recommended to be used in patients affected by hypertension, kidney diseases, liver or cardiovascular disorders or hypokalemia, as they are more sensitive to the adverse effects of licorice (10).
- Patients taking this medication should not take other licorice containing products as serious adverse events may occur such as water retention, hypokalemia, hypertension, cardiac rhythm disorders (10).
- The use is not recommended for children under 4 years of age (11).

9. Interactions with other medicinal products and other forms of interaction

- Not to be used concomitantly with diuretics, cardiac glycosides, corticosteroids, stimulant laxatives or other medications which may aggravate electrolyte imbalance (10).
- The product may counteract antihypertensive action of prescribed medications (7).
- **Azole antifungals:** Licorice may increase the levels of azole antifungals; avoid concurrent use (8).
- Concurrent use with grapefruit juice may increase corticosteroid action of licorice (10).
- **Ciprofloxacin:** The medicinal product may affect the absorption, distribution, and elimination of ciprofloxacin. If the two are used concurrently, their dosages should be separated by at least 2 hours (9).

10. Fertility, pregnancy and lactation

- The use during pregnancy and lactation should be avoided (22).
- No fertility data available.

11. Effects on ability to drive and use machines

No studies on the effect on the ability to drive and use machines have been performed.

12. Undesirable effects



- Allergic reactions affecting the skin, the respiratory system and gastrointestinal tract may occur (10).
- Gastric disorders may occur (11).
- If adverse effects occur, a doctor or a pharmacist should be consulted.

13. Overdose (7, 23)

Prolonged use (more than 4 weeks) and/or intake of high amount of the product, may lead to overdose, which could result in symptoms such as water retention, hypokalemia, hypertension, cardiac rhythm disorders, hypertensive encephalopathy and in rare cases, myoglobinuria.

14. Relevant biological activities

Not required as per Egyptian guidelines for registration of herbal medicines.

15. Additional information

-

16. Date of compilation/last revision

01/06/2026.



References

1.	WHO monographs on selected medicinal plants (2007). Monographs on selected medicinal plants, 3, 42-63.
2.	WHO monographs on selected medicinal plants (2007). Monographs on selected medicinal plants, 3, 136-144.
3.	WHO monographs on selected medicinal plants (1999). Monographs on selected medicinal plants, 1, 183-194.
4.	https://www.gbif.org/species/5341442
5.	https://www.gbif.org/species/7793938
6.	European Union Herbal Monograph on <i>Foeniculum vulgare</i> Miller subsp. <i>vulgare</i> var. <i>vulgare</i> , fructus (2024). Committee on Herbal Medicine Products (HMPC).
7.	Community Herbal Monograph on <i>Glycyrrhiza glabra</i> L. and/or <i>Glycyrrhiza inflata</i> Bat. and/or <i>Glycyrrhiza uralensis</i> Fisch., radix (2012). EMA/HMPC/571119/2010. Committee on Herbal Medicinal Products (HMPC).
8.	Natural Health Product Licorice - <i>Glycyrrhiza glabra</i> (2018). Health Canada, http://webprodhc-sc.gc.ca/nhp/nd-bdipsn/atReq.do?atid=licorice.regliste&lang=eng
9.	Skidmore-Roth, L. (2010). Mosby's Handbook of Herbs & Natural Supplements. 4th ed. ISBN 9780323057417.
10.	Rizzato, G., Scalabrin, E., Radaelli, M., Capodaglio, G. and Piccolo, O. (2017). A new exploration of licorice metabolome. <i>Food Chemistry</i> , 221 , 959–968.
11.	Community Herbal Monograph on <i>Thymus vulgaris</i> L. and <i>Thymus zygis</i> L., Herba. EMA/HMPC/342332/2013. Committee on Herbal Medicinal Products (HMPC).
12.	WHO monographs on selected medicinal plants (2007). Monographs on selected medicinal plants, 3, 259-266
13.	Faudale, M., Viladomat, F., Bastida, J., Poli, F. and Codina, C. (2008). Antioxidant activity and phenolic composition of wild, edible, and medicinal fennel from different Mediterranean countries. <i>J. Agric. Food Chem.</i> 56 , 1912–1920.
14.	Pastorino, G., Cornara, L., Soares, S., Rodrigues, F., and Oliveira, M. (2018). Liquorice (<i>Glycyrrhiza glabra</i>): A phytochemical and pharmacological review. <i>Phytotherapy</i>



Egyptian Herbal Monograph

	<i>Research</i> , 32 (12), 2323–2339. https://doi.org/10.1002/ptr.6178 .
15.	Mamedov, N. A. and Egamberdieva, D. (2019). Phytochemical constituents and pharmacological effects of Licorice: A Review. <i>Plant and Human Health</i> , 3 , 1-21.
16.	Sarfaraz, D., Rahimalek, M. and Saeidi, G. (2021). Polyphenolic and molecular variation in <i>Thymus</i> species using HPLC and SRAP analyses. <i>Sci. Rep.</i> , 11 , 5019.
17.	ESCOP Monographs (2019). Foeniculi fructus, Fennel Fruit. European Scientific Cooperative on Phytotherapy. Simon Mills and Roberta Hutchins, editors. Thieme, Stuttgart. Online series, IBN 978-1-901964-64-6.
18.	Blumenthal, M. (1998). The Complete German Commission E Monographs. Austin, Texas: American Botanical Council.
19.	https://www.rxlist.com/thyme/supplements.htm
20.	PDR for Herbal Medicines (2002). Montvale, NJ: Medical Economics Company, 2 nd ed, ISBN 1-56363-361-2.
21.	Barnes, J., Anderson, L. A. and Phillipson, J. D. (2007). Herbal Medicines, 3 rd edition. Published by the Pharmaceutical Press. ISBN 978 0 85369 623 0.
22.	European Scientific Cooperative on Phytotherapy (ESCOP) (2020). Table of herb-drug interactions based on the monographs https://escop.com/interactions/ Updated: 4/5/2020.
23.	https://www.herbalgram.org/resources/expanded-commission-e/licorice-root/ .