

## The Egyptian Non-prescription Medicinal Products (OTC) List

| No. | Generic name                                       | Dosage form or Route | Notes   |
|-----|--|----------------------|---|
| 1   | Aceclofenac  | Topical              |   |
| 2   | Minerals   | Oral, topical        |   |
| 3   | Vitamin A  | Oral, topical        | <ul style="list-style-type: none"> <li>• Accepted within daily dietary intake in multi-vitamins products,</li> <li>• in case single product with high doses and for specific indication it should be POM</li> </ul> |
| 4   | Vitamin D  | Oral, topical        | <ul style="list-style-type: none"> <li>• Accepted within daily dietary intake in multi-vitamins products,</li> <li>• in case single product with high doses and for specific indication it should be POM</li> </ul> |
| 5   | Vitamin E  | Oral, topical        | <ul style="list-style-type: none"> <li>• Accepted within daily dietary intake in multi-vitamins products,</li> <li>• in case single product with high doses and for specific indication it should be POM</li> </ul> |
| 6   | Vitamin K  | Oral, topical        | <ul style="list-style-type: none"> <li>• Accepted within daily dietary intake in multi-vitamins products,</li> <li>• in case single product with high doses and for specific indication it should be POM</li> </ul> |
| 7   | Vitamin B  | Oral, topical        |   |
| 8   | Vitamin C  | Oral, topical        |   |
| 9   | Aluminium Hydroxide                                | <b>Oral</b>          | • Adequately labeling mentioning impaired renal function precautions  |
| 10  | Ambroxol   | Oral                 |   |
| 11  | Antazoline   | Topical              |   |
| 12  | Aspirin<br>(75,81,100,150,162,250,<br>300,320,325) | Oral                 | • Concentrations above 325mg POM  |
| 13  | Benproperine                                       | Oral                 | • May cause dizziness   |

| No. | Generic name                           | Dosage form or Route                       | Notes  |
|-----|--|--|--|
| 14  | Benzocaine                             | Topical spray, Oral spray & gel & lozenges | <ul style="list-style-type: none"> <li>• <b>Adequate labeling mentioning Methemoglobinemia warning.</b></li> <li>• Used only in the following age groups: <ul style="list-style-type: none"> <li>- Dental gel: From 2 years, only with medical supervisio</li> <li>- Oral solution: From 6 years</li> <li>- Dental spray: From 6 years</li> <li>- Lozenges: From 12 years</li> </ul> </li> </ul> |
| 15  | Bisacodyl                              | Oral, rectal                               |  |
| 16  | Bromhexine                             | Oral                                       |  |
| 17  | Brompheniramine                        | Oral                                       | <ul style="list-style-type: none"> <li>• Use of this drug is completely prohibited in children less than 2 years.</li> <li>• Use of this drug is prohibited in children less than 6 years without medical supervision.</li> </ul>  |
| 18  | Butenafine                             | Topical                                    |  |
| 19  | Caffeine                               | Oral                                       | <ul style="list-style-type: none"> <li>• In combination products only.</li> </ul>  |
| 20  | Calamine                               | Topical                                    |  |
| 21  | Calcium carbonate, Magnesium carbonate | Oral                                       | <ul style="list-style-type: none"> <li>• Caution should be generally exercised in the case of patient with impaired renal function.</li> </ul>   |
| 22  | Camphor                                | Topical                                    |  |
| 23  | Carbinoxamine                          | Oral                                       | <ul style="list-style-type: none"> <li>• Contraindicated in patients less than 2 years.</li> <li>• Antihistamines may cause drowsiness. Use caution when driving or using machine.</li> </ul>  |

| No. | Generic name             | Dosage form or Route | Notes   |
|-----|--------------------------|----------------------|---|
| 24  | Carboxymethylcellulose   | Eye drops            |   |
| 25  | Cetirizine               | Oral                 | <ul style="list-style-type: none"> <li>• Contraindicated in infants less than 6 month.</li> <li>• In chronic conditions, it should be used under medical supervision.</li> <li>• <b>May cause drowsiness; Use caution when driving a motor vehicle or operating machinery</b></li> </ul>  |
| 26  | Chlorhexidine Gluconate  | Topical              |   |
| 27  | Chlorpheniramine Maleate | Oral                 | <ul style="list-style-type: none"> <li>• Use of this drug is completely prohibited in children less than 2 years. Use of this drug is prohibited in children less than 6 years without medical supervision.</li> <li>• <b>May cause drowsiness; caution when driving a motor vehicle or operating machinery.</b></li> <li>• In chronic cases, it must be used under medical supervision.</li> </ul> |
| 28  | Cinchocaine              | Topical              |   |
| 29  | Citric Acid              | Oral                 |   |
| 30  | Clemastine               | Oral                 | <ul style="list-style-type: none"> <li>• Sedation side effects should be clearly labeled.</li> <li>• In chronic conditions, it must be used under medical supervision.</li> </ul>   |
| 31  | Clotrimazole             | Topical              |   |
| 32  | Cromoglicic acid         | Nasal                |   |
| 33  | Cromolyn sodium          | Nasal                | <ul style="list-style-type: none"> <li>• Must not be used in children under 2 years unless directed by a doctor. (to include or to not include age limits)</li> </ul>   |
| 34  | Crotamiton               | Topical              |   |

| No. | Generic name        | Dosage form or Route | Notes   |
|-----|---------------------|----------------------|---|
| 35  | Cyproheptadine      | Oral                 | <ul style="list-style-type: none"> <li>• Sedation side effects should be clearly labeled</li> <li>• Take care in geriatric because of atropine like effects.</li> </ul>   |
| 36  | Desloratadine       | Oral                 | <ul style="list-style-type: none"> <li>• Antihistamines may cause drowsiness. Use caution when driving or using machine.</li> </ul>   |
| 37  | Dexbrompheniramine  | Oral                 | <ul style="list-style-type: none"> <li>• Antihistamines may cause drowsiness. Use caution when driving or using machine.</li> <li>• In case of chronic cases, it must be used under medical supervision.</li> </ul>   |
| 38  | Dexchlorpheniramine | Oral                 | <ul style="list-style-type: none"> <li>• Sedation side effects should be clearly labeled (daily med)</li> <li>• In case of chronic cases, it should be used under medical supervision.</li> </ul>   |
| 39  | Dexpanthenol        | Topical              |   |
| 40  | Diclofenac Sodium   | Topical              |   |
| 41  | Dicyclomine         | Oral                 |   |
| 42  | Dihydroxy Aluminum  | Oral                 | <ul style="list-style-type: none"> <li>• <b>Adequately labeling mentioning impaired renal function precautions</b></li> <li>• Used with extreme caution and only in necessary cases of ascites resulting from liver failure, hypertensive patients and heart diseases.</li> </ul>   |
| 43  | Dimethicon          | Oral & Topical       |   |
| 44  | Dimetindene maleate | Oral & Topical       | <ul style="list-style-type: none"> <li>• Antihistamines may cause drowsiness. Use caution when driving or using machine.</li> <li>• Sedation side effects should be clearly labeled</li> <li>• Not used in children less than 1 year.</li> <li>• Topical anti-histaminic may cause hypersensitivity; avoid in eczema; photosensitivity</li> </ul> <p>Not recommended for longer than 3 days</p> |

| No. | Generic name                   | Dosage form or Route | Notes   |
|-----|--------------------------------|----------------------|---|
| 45  | Diphenhydramine                | Oral & Topical       | <ul style="list-style-type: none"> <li>• Sedation side effects should be clearly labeled.</li> <li>• Topical anti-histaminic may cause hypersensitivity; avoid in eczema; Not recommended for longer than 3 days in photosensitivity</li> <li>• Use of this drug is completely prohibited in children less than 2 years. Use of this drug is prohibited in children less than 6 years without medical supervision.</li> </ul>   |
| 46  | Docosanol                      | Topical              |   |
| 47  | Domiphen bromide               | Oral                 |   |
| 48  | Doxylamine succinate           | Oral                 | <ul style="list-style-type: none"> <li>• In chronic cases, it should be used under medical supervision.</li> <li>• May cause drowsiness. Use caution when driving or using machinery.</li> <li>• Use of this drug is completely prohibited in children less than 2 years. Use of this drug is prohibited in children less than 6 years without medical supervision</li> <li>• Use of this drug in pregnancy is prohibited as it may cause birth defects.</li> </ul>   |
| 49  | Doxylamine succinate, Caffeine | Oral                 | <ul style="list-style-type: none"> <li>• In chronic cases, it should be used under medical supervision.</li> <li>• May cause drowsiness. Use caution when driving or using machinery.</li> <li>• Use of Doxylamine is completely prohibited in children less than 2 years. Use of this drug is prohibited in children less than 6 years without medical supervision</li> <li>• Use of Doxylamine in pregnancy is prohibited as it may cause birth defects.</li> </ul> |
| 50  | Etofenamate                    | Topical              |   |
| 51  | Fexofenadine                   | Oral                 | <ul style="list-style-type: none"> <li>• Not used in children less than 2 years.</li> </ul>   |
| 52  | Folic acid                     | Oral (< 1 mg)        |   |

| No. | Generic name                           | Dosage form or Route           | Notes  |
|-----|--|--------------------------------|--|
| 53  | Guaifenesin                            | Oral & rectal                  | <ul style="list-style-type: none"> <li>Use of this drug is completely prohibited in children less than 2 years. Use of this drug is prohibited in children less than 6 years without medical supervision.</li> </ul> |
| 54  | Hexetidine                             | Mouth Wash                     |  |
| 55  | Hydrocortisone                         | Topical                        | Prolonged use of corticosteroids may increase the incidence of secondary infection.  |
| 56  | Hyoscine                               | Oral                           |  |
| 57  | Ibuprofen                              | Topical                        |  |
| 58  | Indomet(h)acin                         | Topical (maximum strength 1 %) |  |
| 59  | Iron                                   | Oral                           | <ul style="list-style-type: none"> <li>In combinations only</li> <li>ADR: Constipation, darkening of stools, Dental discoloration, Urine discoloration.</li> </ul>   |
| 60  | Lactulose                              | Oral                           |  |
| 61  | Lidocaine                              | Topical                        | Lidocaine 2 % products are not approved to treat teething pain as use in infants and young children can cause serious harm, including death  |
| 62  | Loratadine                             | Oral                           | <ul style="list-style-type: none"> <li>In chronic cases, it should be used under medical supervision.</li> <li>May cause drowsiness. Use caution when driving or using machinery.</li> </ul>                         |
| 63  | lysosyme HCl ,<br>Dequalinium Chloride | Lozenges                       |  |
| 64  | Magaldrate                             | Oral                           |  |
| 65  | Magnesium Hydroxide                    | Oral                           |  |
| 66  | Magnesium Trisilicate                  | Oral                           |  |

| No. | Generic name       | Dosage form or Route  | Notes   |
|-----|--------------------|---|---|
| 67  | Mebevirin          | Oral  |   |
| 68  | Meclizine          | Oral & rectal   | <ul style="list-style-type: none"> <li>The use of this drug is contraindicated in children under 2 years of age, and in children 2-12 years is allowed only under medical supervision.</li> </ul>   |
| 69  | Menthol            | Topical   |   |
| 70  | Mepyramine maleate | Oral & Topical  | <ul style="list-style-type: none"> <li>For the oral dosage forms:                             <ol style="list-style-type: none"> <li>In chronic cases, it should be used under medical supervision.</li> <li>May cause drowsiness. Use caution when driving or using machinery.</li> </ol> </li> </ul>  |
| 71  | Methyl salicylate  | Topical   |   |
| 72  | Miconazole         | Topical & Oral Gel (Maximum strength 2%)                              |   |
| 73  | Minoxidil sulphate | Topical only except gel   | <ul style="list-style-type: none"> <li>Maximum strength 5% for use as OTC</li> </ul>  |
| 74  | Naphazoline        | Nasal (Maximum strength 0.05%) & Ophthalmic (Maximum strength 0.015%) | <ul style="list-style-type: none"> <li>Treatment duration should not exceed 5 days; frequent or prolonged use may cause nasal congestion to recur or worsen. [FDA &amp; technical committee decision]</li> <li>As a single agent it should not be used in children &lt;12 years as a single agent</li> <li>As a combination product with prednisolone it should not be used in children &lt; 15 years.</li> </ul> |
| 75  | Naproxen           | Topical   |   |
| 76  | Nicotine           | Oral Gum  |   |
| 77  | Nonoxynol-9        | Vaginal suppository   | <ul style="list-style-type: none"> <li>Take care of risk of infection (STD) Std: Sexually transmitted disease.</li> </ul>   |
| 78  | Noscapine          | Oral  | <ul style="list-style-type: none"> <li>Maximum single dose 30mg, maximum daily dose 150mg.</li> </ul>   |

| No. | Generic name                           | Dosage form or Route  | Notes  |
|-----|--|---|--|
| 79  | Oral rehydration salts                 | powder for oral solution  |  |
| 80  | Oxetacaine                             | Oral  | <ul style="list-style-type: none"> <li>• Maximum dose 10mg , Maximum daily dose 30mg</li> </ul>  |
| 81  | Oxymetazoline                          | Nasal   | <ul style="list-style-type: none"> <li>• Treatment duration should not exceed 3 days; frequent or prolonged use may cause nasal congestion to recur or worsen.</li> <li>• used for adults and children over 12 years of age</li> <li>• Maximum Strength 0.05% &amp; 0.025%</li> </ul>  |
| 82  | Paracetamol                            | All dosage forms except for: 1.Modified release tablets 2.Prolonged release tablets 3.Injection     | <ul style="list-style-type: none"> <li>• Adequately labeling mentioning hepatitis warning</li> <li>• <b>Maximum dose 1g, maximum daily dose 4g for adult &amp; conc 500mg , 1gm</b></li> </ul>   |
| 83  | Permethrin                             | Topical only<br>1% shampoo Anti-lice<br>2.5% cream , emulsion,<br>lotion ,<br>ointment Anti-scabies | <ul style="list-style-type: none"> <li>• leave on hair for 10 min ( but no longer), use for a single application but may repeat 7 days after first treatment if lice still present ( uptodate)</li> </ul>  |
| 84  | phenylephrine                          | oral  | <ul style="list-style-type: none"> <li>• Not use in children less than 6 years , present in EDA as oral combinations but present as single component eye drops, ear , ophthalmic</li> </ul>  |
| 85  | Pheniramine maleate, Ammonium chloride | Oral  | <ul style="list-style-type: none"> <li>• May cause drowsiness, caution should be taken when driving or operating machinery.</li> </ul>   |
| 86  | Picosal phosphate (Na)                 | Oral <b>drops , tablets</b>   | <ul style="list-style-type: none"> <li>• Short term use only, Maximum dose 7.5mg per day as sodium picosulfate hydrate for adults and adolescents 15 years of age and over. For children between 11 and 14, maximum dose 4.5mg per day. General sale if for adults and children over 10 years. Maximum strength 0.1%, maximum dose 15mg, maximum daily dose 15mg, (AESGP)</li> </ul> |
| 87  | Piperonyl butoxide                     | Topical only  |  |
| 88  | Piroxicam                              | Topical only  | <ul style="list-style-type: none"> <li>• Maximum strength 0.5% in adults and children over 12 years, for a maximum of 7 days. UK. , AESGP</li> </ul>   |



| No. | Generic name                  | Dosage form or Route  | Notes   |
|-----|-------------------------------|---|---|
| 89  | Polyethylene glycol 3350      | Oral  | • Can't be used for more than 7 consecutive days, Indicated in temporary constipation in adults and children over 6 months.(france AESGP)   |
| 90  | Polymyxin B                   | Topical only  |   |
| 91  | Povidone-Iodine               | Topical only  |   |
| 92  | Pramocaine                    | Topical only  |   |
| 93  | Promethazine                  | Oral  | • Use is completely prohibited in children < 2 years, Use in children <6 years must only be under medical supervision   |
| 94  | Propanthelin                  | Topical only  |   |
| 95  | Pseudoephedrine               | Oral only   | - 30 mg/5ml:<br>Contraindicated in children less than 12 years old.<br>-15 mg/5ml:<br>Contraindicated in pediatrics less than 2 years old and from 2-12 years old used under medical supervision. |
| 96  | Psyllium Muciloid (Ispaghula) | Oral  | • Administration precautions should be adequately labeled   |
| 97  | pyrethrins                    | Topical   |   |
| 98  | Racecadotril                  | Oral , Powder,hard gelatin capsules,sachets(granules for oral suspension) |   |
| 99  | Selenium                      | Oral tablets  | • Only in Combination   |
| 100 | Simethicon                    | Oral(Chewable tablets ,suspension,oral drops)                             |   |

| No. | Generic name   | Dosage form or Route                                      | Notes  |
|-----|--|---|--|
| 101 | Sodium Bicarbonate   | Oral  | <p><b>Only in Combination</b></p> <p>مدلا طغض بضر م لثم ( حلملا ) مويو صلا مادختسا نم للاقلا مهتم<br/>بولطملا بضر ممل رذحب مدختسي نع جتانلا ءاقستسلا او يولكلا<br/>لشفلاو بلقلا بضر مو<br/>يدبكلا لشفلا</p>  |
| 102 | Sodium Chloride  | nasal spray ,nasal gel & ophthalmic solution as eye drops |  |
| 103 | Sodium Chromoglicate   | Eye drops   | <ul style="list-style-type: none"> <li>• <b>Maximum strength 2%</b></li> <li>• OTC indication: only for the relief and treatment of seasonal and perennial allergic conjunctivitis</li> </ul>  |
| 104 | Terbinafine  | Topical ( Cream,gel,topical solution, spray)              |  |
| 105 | Triamcinolne oral base   | Oral paste  | <ul style="list-style-type: none"> <li>• For painful ulcer.</li> <li>• OTC only in those aged 18 years and over.</li> </ul>  |
| 106 | Tribneoside  | Topical   |  |
| 107 | Tripolidine  | Oral(Tablets ,syrup)                                      | <ul style="list-style-type: none"> <li>• Contraindicated in children less than 2 years.</li> <li>• Must be used under medical supervision in children less than 6 years.</li> <li>• Must be used under medical supervision in chronic cases.</li> <li>• Antihistamines may cause drowsiness. Use caution when driving or using machinery.</li> <li>• Take care in geriatric because of atropine like effects.</li> </ul> |
| 108 | Xylometazoline   | Nasal drops   | <ul style="list-style-type: none"> <li>• <b>Maximum strength : up to 0.1%</b></li> <li>• Prohibited in children &lt; 6 years, and</li> <li>• pharmacology warning.</li> </ul>  |
| 109 | Zinc Sulphate  | Oral  |  |
| 110 | Miconazole Nitrate 2%<br>Hydrocortisone<br>Micronized 1% Cream | Topical   |  |

| No. | Generic name  | Dosage form or Route  | Notes  |
|-----|---|-----------------------|--|
| 111 | Fexofenadine Hydrochloride 60mg +Pseudoephedrine Hydrochloride 120mg                    | Oral                  | • Used only in those aged 2 years and over.  |
| 112 | Bacillus Clausii spores 2 Billion/5ml   | Oral                  |  |
| 113 | Paracetamol 500 mg Caffeine 65mg  | Oral                  |  |
| 114 | Sodium alginate 500mg/10ml, Sodium Bicarbonate 267mg/10ml, Calcium Carbonate 160mg/10ml | Oral sachet           | • Use with caution in patients on sodium-restricted diet for example in patients with hypertension, heart disease, renal failure and ascites from hepatic failure. |
| 115 | Sodium alginate 500mg/5ml, Sodium Bicarbonate 100mg/5ml                                 | Oral suspension       | • Use with caution in patients on sodium-restricted diet for example in patients with hypertension, heart disease, renal failure and ascites from hepatic failure. |
| 116 | Sodium alginate 250mg, Sodium Hydrogen carbonate 133.5, Calcium carbonate 80mg          | Oral Chewable Tablets | • Use with caution in patients on sodium-restricted diet for example in patients with hypertension, heart disease, renal failure and ascites from hepatic failure. |
| 117 | Beta-Sitosterol 0.25g/100g  | Topical ointment      |  |

**List of Newly OTC switch-approved products According to Guidelines for Classification as Nonprescription Medicinal Products (OTC)**

**September 2023**

|   | <b>Trade Name</b>                            | <b>Active Ingredient</b>   | <b>Approved Pack Limit</b>   |
|---|--|--|--|
| 1 | <b>Levohistam Film Coated Tablets</b>        | Levocetirizine Dihydrochloride 5mg                                   | Carton box contain 1, 2 or 3 (Al/PVC) strips, each of 10 film coated tablets + insert leaflet  |
| 2 | <b>Levohistam 5mg/10ml Syrup</b>             | Levocetirizine Dihydrochloride 5mg /10ml                             | Carton box containing 135ml amber PET palstic bottle contains 120 ml solution with HDPE plastic closure with compressible (LDPE) plastic foam liner and red tamper evident ring and 20 ml measured plastic cup with insert leaflets.   |
| 3 | <b>Levohistam 5mg/1ml oral drops</b>         | Levocetirizine Dihydrochloride 5mg /1ml                              | Carton box contains amber (Type III) sodalime-silica glass bottle containing 10 ml or 20 ml solution with white (HDPE) plastic pilfer proof cap, (LDPE) internal foam liner and attached with white LDPE plastic 1ml dropper attached to HDPE plastic nozzle with an insert leaflet  |
| 4 | <b>Levcet Melt in mouth tablets</b>          | Levocetirizine Dihydrochloride 5 mg                                  | Carton box containing 1 , 2 (Al/Opaque PVC / PVDC ) strip each of 10 tablets & an inner leaflet .  |
| 5 | <b>Levcet Oral Solution</b>                  | Each 5 ml contains:<br>Levocetirizine Dihydrochloride 2.5 mg         | Carton box containing amber PET (Polyethylene terephthalate ) bottle containing 120 ml ( capacity $149.6 \pm 3$ ml ) oral solution with white polypropylene closure cap containing white solid PE foam ( low density polyethylene and carbon dioxide ) inside with graduated transparent polypropylene plastic measuring cup and an insert leaflet . |
| 6 | <b>Antopral 20 mg</b>                        | Pantoprazole sesquihydrate 22.55 mg equivalent to Pantoprazole 20 mg | Carton box containing 2 strips (Al/Al) each of 7 gastro-resistant tablets and inner leaflet.   |
| 7 | <b>Rani-F Effervescent Granules</b>          | Famotidine 0.02 gram /sachet   | Carton box containing 6,12 or 60 aluminum laminated foil sachets each sachet consists of 4 layers from inside to outside (LDPE/Aluminium/LDPE/paper), each sachet contains 5 gm effervescent granules with insert leaflet.   |
| 8 | <b>Solofresh sterile Ophthalmic Solution</b> | Sodium Hyaluronate 2 mg / ml   | Carton box containing (LDPE ) plastic bottle containing 10 ml solution with white LDPE plastic self-dropper covered with white PP plastic cover + insert leaflet   |

|    | <b>Trade Name</b>   | <b>Active Ingredient</b>   | <b>Approved Pack Limit</b>   |
|----|---|--|--|
| 9  | <b>Nasacort AQ<br/>Nasal Spray<br/>Aqueous<br/>Suspension</b> | Triamcinolone Acetonide 55mcg<br>(0.055% w/w or mg/actuation)  | A carton box containing a 20ml high density polyethylene (HDPE) bottle fitted with a metered-dose spray pump unit, which contains 16.5gm suspension (with 9.075mg triamcinolone acetonide) providing 120 actuation & an inner leaflet.   |
| 10 | <b>Maalox Control<br/>Gastro-resistant<br/>tablets</b>        | Pantoprazole 20 mg as sodium sesquihydrate 22.575 mg   | Carton box contains one (Alu/Alu) blister of 14 gastro-resistant tablets with insert leaflet.  |
| 11 | <b>Enterogermina 4<br/>billion / 5 ml</b>                     | Each 5 ml vial contains :<br>Spores of poly-antibiotic resistant B.clausii 4 billion                               | Cardbord cartons containing 10 low density polyethylene bottles produced by plastic extrusion each containing 5 ml oral suspension and inner leaflet .   |
| 12 | <b>Essential Capsule<br/>300 mg</b>                           | Phospholipids from soya Beans 76% (3 sn-phosphatidyl )choline 300 mg   | Carton box containing 3 Aluminum / Perlalux Tristar ultra (Transparent PVC /PE / PVDC ) blisters<br>Each of 10 H.G.C. +Inner leaflet.  |
| 13 | <b>Melacryst Oro-<br/>dispersible film</b>                    | Melatonin 1.5 mg   | Carton box containing 10,20,30 Oro-dispersible films ,each one film packed in a four layer pouch form outside to inside (paper ,LDPE , Aluminium ,LDPE ) with insert leaflet.  |
| 14 | <b>Hemoclar topical<br/>cream</b>                             | Each 1 gram contains :<br>Pentosan sulphuric polyester 5 mg  | Carton box containing printed sealed collapsible aluminium tube of 40 gm cream closed with HDPE cap with insert leaflet.   |
| 15 | <b>Brufen 200 mg<br/>Sugar Coated<br/>tablet</b>              | Ibuprofen 200 mg   | Carton box containing 3 (Al/transparent PVCs )strips, each of 10 ssugar-coatedtablets and an insert leaflet.   |
| 16 | <b>Brufen syrup</b>   | Ibuprofen 100mg/5ml Syrup  | Carton box containing an amber glass (type III ) bottle of 110 or 150 ml syrup and an insert leaflet.  |
| 17 | <b>Brimillergy<br/>Ophthalmic<br/>solution</b>                | Each 1 ml solution contains :<br>Brimonidine Tartrate 0.25 mg  | Carton box containing 10 ml white plastic bottle (LDPE ) contains 7.5 ml solution with (LDPE ) dropper and white (PP ) plastic cover with insert leaflet .   |
| 18 | <b>Notussil Oral<br/>Suspension</b>                           | Each 5 ml of oral suspension contains :<br>Cloperastine Fendizoate 35.4 mg<br>equivalent to Cloperastine HCl 20 mg | -Carton box containing polyethylene terephthlate amber glass (Type III ) bottle (its capacity $79.7 \pm 2$ ml ) containing 60 ml and pamphlet .<br>- Carton box containing polyethylene terephthlate amber glass (Type III ) bottle ( its capacity $140 \text{ ml} \pm 2.8$ ml ) containing 120 ml and pamphlet. |

|    | Trade Name   | Active Ingredient  | Approved Pack Limit  |
|----|--|--|--|
| 19 | <b>Gastrazole Capsules</b>                         | Each 240 mg pellets contain :<br>Omeprazole 20 mg  | Carton box contains HDPE plastic bottle lined with (Al / transparent colorless LDPE ) from outside to inside induction seal containing 14 hard gelatin capsules and LDPE white plastic capsules containing white silica gel desiccant covered with protective paper layers with HDPE outer plastic screw cap and HDPE inner plastic cap lined with protective layers of paper with insert leaflet. |
| 20 | <b>Evastine Film Coated Tablets</b>                | Ebastine 10 mg   | Carton box containing 1 or 2 ( Al / transparent PVC ) strips , each of 10 film coated tablets with insert leaflets.  |
| 21 | <b>Nizapex Topical Shampoo</b>                     | Each 1 gm contains : Ketoconazole 20 mg  | Carton box containing a plastic ( shoulder material : HDPE & sleeve material : LDPE ) tube of 80 ml shampoo & closed with polypropylene cap with insert leaflet.   |
| 22 | <b>Ticanase Nasal spray</b>                        | Each actuation contains : Fluticasone Propionate 50 mcg  | Carton box containing amber glass type (III) bottle of capacity 15 ml containing 7 gm suspension with white polypropylene nasal sprayer head delivers 60 puffs ,capped with transparent polypropylene cap and INOX pump with insert leaflet.   |
| 23 | <b>Movicol Powder for oral solution in sachets</b> | Each 13.8 gm sachet contains :<br>-Macrogol 3350 13.125 gm<br>-Sodium Bicarbonate 0.1785 gm<br>-Sodium Chloride 0.3507 gm<br>-Potassium Chloride 0.0466 gm | Carton box containing 20 sachets ,each of 13.8 gm powder made of laminate consisting of four layers (Paper /LPDE / LA /LDPE ) assembled in pairs duplex separated by perforation line with insert leaflet .  |
| 24 | <b>Radian Massage cream Topical Cream</b>          | Camphor 1.410% W/W + Camphor oil white 0.320% W/W + Capsicum Oleoresin 0.005% W/W + Menthol 2.540% W/W +Methylsalicylate 0.420% W/W                        | Carton box containing lacquered aluminum tube with aluminum seal of 40gm or 100gm cream with (HDPE) screw cap inner leaflet.   |
| 25 | <b>Bronchicum Elixir</b>                           | Each 5 ml contains:<br>Thyme fluid extract 0.325 g<br>Primula root fluid Extract 0.162 g   |  |
| 26 | <b>Movicol Pediatric Plain</b>                     | Each 6.9 gm sachet contains:<br>-Macrogol 3350 6.563 gm<br>-Sodium Chloride 175.4 mg<br>-Sodium Hydrogen Carbonate 89.3 mg<br>-Potassium Chloride 25.1 mg  | Carton box containing 30 sachets, each of 6.9 gm powder made of laminate consisting of four layers (paper/LDPE/AL foil/LDPE) with insert leaflet.  |

|    | Trade Name                                       | Active Ingredient   | Approved Pack Limit   |
|----|--|---|---|
| 27 | <b>Stomatect Oral suspension</b>                 | Each 5 ml contains :<br>Potassium Bicarbonate 100 mg<br>Sodium Alginate 500 mg  | Carton box containing amber glass type III containing 150 or 300 ml suspension with white circular polypropylene cap havinf LDPE red ring at the edge and contain a foam PVDC stopper at upside with insert leaflet . |
| 28 | <b>Sinupret Coated Tablets</b>                   | Gentian root Gentiana Radix 6 mg ,Primerose flowers (Primulae flows ) with calyx 18 mg ,Common sorrel herb (Rumicis Herba ) 18 mg , Elder flowers ( Sambuci flows ) 18 mg ,Shop vervain wort (Verbenae herba ) 18 mg                                | Carton box containing (Al/ Transparent PVC / PVDC ) strips ,each of 25 coated tablets and inner leaflet.  |
| 29 | <b>Stopadol Night Film Coated Tablets</b>        | Paracetamol 500 mg +<br>Diphenhydramine Hydrochloride 25 mg   | Carton box containing 1 , 2 or 3 (Aluminum , transparent PVC ) strips , each of 10 film coated tablets with insert leaflet .  |
| 30 | <b>Pentamix Syrup</b>                            | Each 5 ml contains:<br>-Tilia flowers aqueous extract 0.0625 gm<br>-Guava Leaves aqueous extract 0.125 gm<br>-Bitter Fennel fruits aqueous extract 0.05 gm<br>-Eucalyptus leaves aqueous extract 0.025 gm<br>-Thymus leaves aqueous extract 0.05 gm | Carton box contains amber glass bottle (type II) with white plastic cap with polyethylene foam and inner leaflet.   |
| 31 | <b>Nercafix Oro-dispersible Film</b>             | Caffeine anhydrous 40 mg  | Carton box contains 10 or 20 or 30 oral dispersible films , each film packed in four layer pouch , from outside to inside (Paper , LDPE ,Aluminum & LDPE ) with insert leaflet.                                       |
| 32 | <b>Algason Massage Cream</b>                     | Each 1 g contains: Diethylamine Salicylate 0.1 gm+ Camphor 0.014 gm +Menthol 0.025 gm   | Carton box inside it printed aluminum tube of 40 g cream with Al seal and white (HDPE) cap with insert leaflet.   |
| 33 | <b>Glycerin for adults Rectal suppositories</b>  | Glycerin 1.4 gm   | Carton box containing one white opaque (PVC / LDPE) strip containing 5 suppositories with insert leaflet .  |
| 34 | <b>Glycerin for infants Rectal suppositories</b> | Glycerin 0.7 gm   | Carton box containing one white opaque ( PVC / LDPE) strip containing 5 suppositories with insert leaflet.  |
| 35 | <b>Miopan Plus Oral suspension</b>               | Each 5 ml suspension contains:<br>Magaldrate (Anhydrous) 0.54 gm<br>Simethicone 0.04 gm   | Carton box containing amber brown plastic (PET) bottle containing 100 ml suspension with HDPE plastic cap with insert leaflet.  |



|    | Trade Name   | Active Ingredient   | Approved Pack Limit   |
|----|--|---|---|
| 36 | <b>Digicomag Chewable Tablets</b>                            | Calcium Carbonate<br>680 mg<br>Simethicone 25 mg<br>Magnesium Carbonate ( Heavy )<br>80 mg  | Carton box containing 3 (AL/Transparent PVDC) blisters each contains 10 chewable tablets with insert leaflet.   |
| 37 | <b>Moov massage Topical cream</b>                            | Each 1 g contains:<br>Oleoresin capsicum 35% 0.42mg<br>Camphor 12 mg<br>Menthol 12.1mg<br>Methyl salicylate 4.8 mg<br>Oil of camphor 1.8 mg   | Carton box containing AL laminated tube of 40gm or 100gm cream with (HDPE) screw cap and inner leaflet  |
| 38 | <b>Chitocal Hard Gelatin Capsules</b>                        | Chitosan Powder<br>500 mg<br>(Standardized deacetylation degree NLT 70% and NMT 90%)<br><br>*Gymnema Sylvester leaves powder extract 50 mg<br>(Standardized to 25% Gymnemic acid )<br><br>*Vitamin C ( ascorbic acid)<br>100 mg | Carton box containing 2,3,6 or 8 (Al / transparent PVC) strips each of 10 H.G.C and an inner leaflet.   |
| 39 | <b>Quick – Slim Hard Gelatin Capsules Containing Pellets</b> | Each Capsule contains 240 mg of 50% Orlistat Pellets containing:<br>Orlistat 120 mg   | Carton box contains 1 or 3 (Al / transparent colorless PVC ) strips of 10 hard gelatin capsules with insert leaflet.  |
| 40 | <b>Emla Topical Cream</b>                                    | Each 1 gm contains:<br>Lidocaine 25 mg<br>Prilocaine 25 mg  | Carton box containing 5 tubes ,each of 5 gm cream ,each tube is a collapsible aluminum tube closed with membrane ,internally coated with epoxy resine protective lacquer ,with polypropylene cap with spike for perforation of the membrane .With insert leaflet. |
| 41 | <b>Ivyrospan Syrup</b>                                       | Each 5 ml contains:<br>Dried Ivy leaf extract 35 mg<br>Standardized as hederacoside C (50 mg/g-250 mg/g )   | Carton box containing amber (type III ) glass bottle containing 100 ml syrup with white polypropylene cap containing white solid IDPE & Carbon dioxide foam with safety seal and transparent ( HDPE) plastic measuring cup & inner leaflet.                       |
| 42 | <b>Agnucaston Film Coated Tablets</b>                        | Dry Extract from Agnus castus fruit<br>20 mg<br>Assay Herbal preparation : Casticin ,<br>0.05 – 0.25 [% ( m / m ) ]   | Carton box containing 3 or 6 (Al / transparent PVC / PVDC) strips , each of 10 film coated tablets + inner leaflet.   |



|    | <b>Trade Name</b>  | <b>Active Ingredient</b>   | <b>Approved Pack Limit</b>  |
|----|--|--|---|
| 43 | <b>Klimadynon®<br/>Film Coated<br/>Tablets</b>                     | Dry extract of Cimcifuga racemosa<br>2.8 mg  | Carton box containing 1, 2 or 3 (Al / colorless PVC / PVDC) strips ,each of 10 film coated tablets + inner leaflet.   |
| 44 | <b>Imoflora Tablet</b>   | Loperamide Hydrochloride 2 mg<br>Simethicone 125 mg  | Carton box containing one ( AL / opaque PVC ) strip of 10 tablets and an inner leaflet.   |
| 45 | <b>Norhinose Nasal<br/>Spray</b>                                   | Each metered dose contains:<br>Mometasone Furoate Monohydrate<br>0.052 mg equivalent to Mometasone<br>Furoate 0.05 mg.   | Carton box containing an amber glass type (III ) 15 ml capacity glass bottle containing 9.7 gm suspension of 60 doses or 18 gm suspension of 120 doses with a white polypropylene plastic nasal sprayer capped with lower density polyethylene (LDPE ) dip tube surrounded by polypropylene housing cap capped with LDPE plastic transparent cap and an insert leaflet. |
| 46 | <b>Algesal Suractive<br/>Topical Cream</b>                         | Each 100 g contains:<br>Diethylamine Salicylate 10 g<br>Myrtecaine (nopoxamine) 1 g                                      | Carton box contains collapsible AL tube internally coated with epoxy and melamine film varnished the mastic layer on the bottom twice varnish with araldite lithographic pattern cylindrical part containing 40 g homogenous cream closed with "HDPE" screw cap + insert leaflet.   |
| 47 | <b>Supermag B6<br/>Film Coated<br/>Tablets</b>                     | Magnesium lactate Dihydrate 470 mg<br>equivalent to<br>Magnesium 48 mg<br>-Pyridoxin Hydrochloride ( Vitamin<br>B6) 5 mg | Carton box contains 1 , 2 or 3 ( Al / transparent (PVC / PVDC) blister , each of 10 film coated tablets with insert leaflet.  |
| 48 | <b>Megafen Tablets</b>   | Ibuprofen 200 mg<br>Paracetamol 325 mg   | Carton box containing 1 ,2 or 3 (Al / transparent PVC ) blisters , each of 10 tablets + an insert leaflet.  |
| 49 | <b>Orly 60 mg Hard<br/>Gelatin capsules<br/>containing pellets</b> | Each hard gelatin capsule contains 120 mg orlistat pellets 50%,equivalent to:<br>Orlistat 60 mg                          | Carton box containing 1 ,2 , 3 , 4 , 5 or 6 (AL / AL ) strips ,each of 10 hard gelatin capsules with insert leaflet.  |
| 50 | <b>Nasonex 50 mcg /<br/>actuation Nasal<br/>spray suspension</b>   | Each gm contains:<br>Mometasone Furoate ( as<br>monohydrate) 0.5 mg  | Carton box containing 23.5 ml white HDPE bottle containing 18 g suspension (140 actuations ) supplied with a metered dose polypropylene spray pump actuator and an inner leaflet  |

|    | Trade Name                                      | Active Ingredient   | Approved Pack Limit   |
|----|---|---|---|
| 51 | <b>Privacond Ophthalmic Solution</b>            | Each 1 ml ophthalmic solution contains:<br>Tetrahydrozoline Hydrochloride 0.5 mg<br>Polyvinyl Alcohol 5 mg<br>Povidone mg | Carton box containing 15 ml ophthalmic solution in white opaque LDPE plastic dropper bottle with white opaque LDPE dropper tip and HDPE screw cap and an inner leaflet.   |
| 52 | <b>Sinupret ® forte Coated Tablets</b>          | Herbal  | Carton box contains 50 coated tablets in blisters , the blisters are made of transparent PVC / PVDC and aluminum foil , each blister contains 25 coated tablets with insert leaflet.  |
| 53 | <b>Avamys Nasal Spray Nasal Spray</b>           | Each spray actuation delivers:<br>Fluticasone Furoate 27.5 mcg  | Carton box contains (off-white plastic device with a dose indicator window , light blue side actuated lever and lid which contains a stopper .The plastic device contains the nasal spray suspension within a Type I amber bottle (glass) fitted with a metering spray pump ) of 120 sprays and inner leaflet.                              |
| 54 | <b>Glycerin Adult Suppositories</b>             | Glycerin 1.47 g   | Carton box contains 2 (LDPE / White Opaque PVC) strips, each strip of 5 suppositories + an inner leaflet.   |
| 55 | <b>Glycerin Infantile Suppositories</b>         | Glycerin 0.7359 g   | Carton box containing 2 ( PVC / PE ) strips , each of 5 suppositories and an inner leaflet.   |
| 56 | <b>Cetafen 200 / 500 mg Film Coated Tablets</b> | Ibuprofen 200 mg<br>Paracetamol 500 mg  | Carton box containing 1 , 2 or 3 (AL / opaque PVC / PVDC ) blisters , each of 10 tablets + inner leaflet.   |
| 57 | <b>Melacryst 3 mg Oro-dispersible Film</b>      | Melatonin 3 mg  | Carton box contains 10 ,20 or 30 Oro-dispersible films ,each is packed in a four-layer pouch (from outside to inside : paper , LDPE , aluminum , LDPE) and insert leaflet.  |
| 58 | <b>Flixonase Nasal spray</b>                    | Each actuation (100 mg) contains:<br>Fluticasone Propionate micronized 0.05 mg  | Carton box containing Amber (type III) glass bottle (weight of contents for 120 doses is not less than 15 gm) fitted with a metering atomizing spray pump with a low density polyethylene dip tube fitted to the pump is a combined polypropylene nasal white actuator covered with a green polypropylene dust cover and an insert leaflet. |

|    | Trade Name   | Active Ingredient  | Approved Pack Limit  |
|----|--|--|--|
| 59 | <b>Agiolax Granules in sachets</b>   | Plantago Ovata seed (Plantago Ovata) 2.6 gm+ Ispaghulla husks(Plantago Ovata Forssk) 0.11 gm+Tinnevelly senna pods (Cassia angustifolia Vahl) 0.34 – 0.66 gm | Carton box containing 12 sachets , each of 5 gm and an insert leaflet.   |
| 60 | <b>Panadol Cold + Flu Vapour Release + Decongestant Powder for Oral Solution</b> | Each sachet contains:<br>Paracetamol 600 mg<br>Phenylephrine Hydrochloride 10 mg   | Carton box contains 10 ( paper ,polyethylene , aluminium foil ,ethylene-methacrylic acid , co polymer (EMAA) laminate sachets with insert leaflet.     |
| 61 | <b>Panadol Sinus Relief PE Tablets</b>   | Paracetamol 500 mg<br>Phenylephrine Hydrochloride 5 mg   | Carton box containing 2 ( Al/Transparent colorless PVDC) strips, each of 12 tablets+ an inner leaflet.   |
| 62 | <b>Contractubex Topical Gel</b>  | Each 100 gm gel contains:<br>Allantoin 1 gm<br>Heparin Sodium 5000 I.U. equivalent to 0.04 gm<br>Cepae fluid extract 10 gm                                   | Carton box contains Al tube contains 20 , 30 , 50 gm & 100 gm gel with aluminum membrane and white screw HDPE plastic cap with insert leaflet          |
| 63 | <b>Menobrocond Film Coated Caplets</b>   | Acetaminophen 500 mg<br>Pamabrom 25 mg   | Carton box containing 1 , 2 or 3 ( Al / opaque white PVC) strips , each of 10 film coated caplets + insert leaflet.                                    |
| 64 | <b>Cetafen Plus N Film Coated Tablets</b>  | Paracetamol 500 mg<br>Ibuprofen 200 mg<br>Caffeine Anhydrous 65 mg   | Carton box containing 1 , 2 or 3 (opaque white PVC - PVDC - AL) strips , each of 10 tablets with insert leaflet.                                       |
| 65 | <b>Flu-C Powder for oral solution in sachets</b>                                 | Each sachet contains:<br>Chloropheniramine Maleate 4 mg<br>Paracetamol (microfined) 500 mg<br>Caffeine Anhydrous 30 mg<br>Ascorbic Acid (Vitamin C) 500 mg   | Carton box containing 6 sachets (from outside to inside paper / LDPE/AL/LDPE) each containing 3 gm powder and insert leaflet.                          |
| 66 | <b>Spasmomen Film Coated Tablets</b>   | Otilonium Bromide 40 mg  | Carton box contains 2 ( Al / white opaque PVC) strips , each of 10 tablets with insert leaflet.  |
| 67 | <b>Fastum 2.5% gel Topical Gel</b>   | Ketoprofen 2.5 g / 100 g   | Carton box containing one tube of 15 ,50 ,60 , 100 gm gel and inner leaflet.   |
| 68 | <b>Purgolaxin Enema</b>  | Each 5 ml solution contains:<br>Sodium Citrate 450 mg<br>Sodium Lauryl Sulfoacetate 45 mg  | Carton box contains 4 single dose white (LDPE) plastic tube with white (LDPE) plastic nozzle each containing 5 ml rectal solution with insert leaflet. |

|    | Trade Name  | Active Ingredient   | Approved Pack Limit  |
|----|---|---|--|
| 69 | <b>Ruscap hard gelatin capsule</b>                    | Ruscus Aculeatus extract 150 mg+<br>Trimethyl hesperidin 150 mg+<br>Ascorbic acid 100 mg  | Carton box containing 1,2 or 3 (AL/Transparent PVC )strips each of 10 hard gelatin capsule & insert leaflet  |
| 70 | <b>Alverinospasm soft gelatin capsule</b>             | Alverine Citrate 60 mg + Simethicone 300 mg   | Carton box containing 1,2 or 3 (AL/PVC) strips each of 12 soft gelatin capsules + insert leaflet.  |
| 71 | <b>Marpalene 0.1% Topical cream</b>                   | Adapalene 1mg in each 1gm   | Carton box containing collapsible Aluminum tube of 30gm cream with white HDPE plastic cap and insert leaflet.  |
| 72 | <b>Vitamax Plus Soft Gelatin Capsules</b>             | Herbal medicine containing multivitamins and herbs  | Carton box containing 1 , 2 or 3 (Al/PVC) strips ,each of 10 soft gelatin capsules and an inner leaflet.   |
| 73 | <b>Expaintech-PM Tablets</b>                          | Paracetamol 500 mg<br>Diphenhydramine Citrate 38 mg   | Carton box containing 1 (AL/ white opaque PVDC) strip of 10 tablets + inner leaflet .  |
| 74 | <b>Anaseziago Topical Cream</b>                       | Each 1 gm contains:<br>Lidocaine 25 mg<br>Prilocaine 25 mg  | Carton box containing one AL tube of 30 gm and an inner leaflet.   |
| 75 | <b>Panadol Cold &amp; Flu Day Film Coated Tablets</b> | Paracetamol 500 mg<br>Caffeine 25 mg<br>Phenylephrine Hydrochloride 5 mg  | Carton box containing 2 (Al+Child resistant foil /Opaque white PVC ) strips , each of 12 film coated tablets and an insert leaflet.                                    |
| 76 | <b>Tussigreen Syrup</b>                               | Each 5 ml Contains:<br>Diphenhydramine Hydrochloride 14 mg<br>Ammonium Chloride 135 mg  | Carton box containing amber brown (Type III) glass bottle containing 100 ml syrup ,with high density polyethylene cap and(LDPE) plastic foam liner and insert leaflet. |
| 77 | <b>Daflon 500 mg Film Coated Tablets</b>              | Each tablet contains: purified, micronized flavonoid fraction, composed of:<br>Diosmin 90% (450 mg)<br>Flavonoids (as hesperidin) 10% (50 mg) | Carton box containing 1 or 2 blisters pf (Transparent PVC / Al) ,each of 15 film coated tablet and inner leaflet.  |
| 78 | <b>Daflon 1000 mg Film Coated Tablets</b>             | Purified, Micronized Flavonoid Fraction1000 composed of:<br>Diosmin 90% (900 mg)<br>Flavonoids (as hesperidin) 10% (100 mg)                   | Carton box contains heat sealed 3 (Al / Transparent PVC ) blisters ,each contains 10 film coated tablets with insert leaflet.  |
| 79 | <b>Controloc 20 mg Gastro resistant Tablet</b>        | Pantoprazole Sodium sesquihydrate 22.57 mg<br>Eq. to Pantoprazole 20 mg   | Carton box contains 1 or 2 (Al/Al) blisters each of 7 tablets and insert leaflet.  |
| 80 | <b>Jeparilon Chewable Tablets</b>                     | Calcium Carbonate 800 mg<br>Famotidine 10 mg<br>Magnesium Hydroxide 165 mg  | Carton box containing 2(Al/Al) strips , each of 5 chewable tablets and insert leaflet.   |

|    | Trade Name                    | Active Ingredient   | Approved Pack Limit  |
|----|-------------------------------|---|--|
| 81 | Stopspasm Film Coated Tablets | Phloroglucinol 62.233 mg<br>Trimethylphloroglucinol 80 mg | Carton box containing 1 , 2 or 3 (Al / PVC ) strips , each strip contains 10 film coated tablets and an inner leaflet. |

- يرجى ملاحظة أنه يتم تغيير وضع المستحضر وإلغاء ادراجه بقائمة الأدوية غير الوصفية ؛ حال ظهور أى معلومات جديدة خاصة بأمنية المستحضر مما يجعله غير ملائماً للاستخدام بواسطة المريض دون اشراف طبي .