

Egyptian Herbal Monograph

Volume 2

Medicinal plants used in Egypt

Egyptian Drug Authority (EDA)

2026





Egyptian Herbal Monograph

Medicinal Plants Used in Egypt

***Grindelia hirsutula* Hook. & Arn**

جرنديليا

1. Names & Synonyms (1)

***Grindelia hirsutula* Hook. & Arn.**

Syns.: *Grindelia robusta* Nutt., *Grindelia humilis* Hook. et Arn., *Grindelia camporum* Greene.

Family: Asteraceae (Compositae)

Arabic: Grindelia جرنديليا

English name: Gumweed (2,3), Gum plant (4).

2. Parts used for medicinal purpose

Herb/ leaves and flowering tops (2-6).

3. Major chemical constituents

- **Resin (labdane type of diterpene acids):** Grindelic acid and its derivatives (7).
- **Acetylenic compounds:** Matricarianol and marticarianol acetate (8).
- **Essential oil:** Mono terpenoids as: α -pinene, β -pinene, borneol, trans-pinocarveol, bornyl acetate and limonene (9-12).
- **Flavonoids:** Mainly quercetin-methylethers (13), beside kaempferol-3-methylether and kaempferol-3,7-dimethylether (8).
- **Others:** Phenolic acids (14), saponins, triterpenoid and tannins (8).

4. Medicinal Uses (Indications)

- Relief of cough associated with cold, catarrh of the upper respiratory tract and bronchitis. (2,5,6).

5. Herbal preparations correlated to medicinal use (2,5)

1. Comminuted herbal substances are added to water as herbal tea in the form of infusion.
2. Liquid extract (1:1), extraction solvent ethanol (22.5% V/V).

3. Tincture (1:10 or 1:5), extraction solvent ethanol (60-80% V/V).

Herbal preparations (2-3) are in liquid dosage forms for oral use. The pharmaceutical form should be described by the pharmacopoeia full standard term.

6. Posology and method of administration correlated to medicinal use

Preparation 1

- 2 - 3 g of the comminuted herbal substance in 150 ml of boiling water as a herbal infusion, up to 3 times daily (2).
- 4 - 6 g of the comminuted herbal substance as a 10% infusion, daily (5).

Preparation 2

- 0.6 - 1.2 ml, 3 times daily. Daily dose: 1.8 - 3.6 ml (2).
- 3 - 6 g of fluid extract, daily (5).

Preparation 3

- 0.5 - 1 ml, 3 times daily. Daily dose: 1.5 - 3 ml (2,5).

Duration of use

If the symptoms persist longer than 1 week during the use of the medicinal product, a doctor or a pharmacist should be consulted (2).

Method of administration: Oral use (2,5).

7. Contraindications (2,5)

- Hypersensitivity to the active substance and to other plants of the same family.

8. Special warnings and precautions for use (2,5)

- If the symptoms worsen during the use of the medicinal product or if dyspnoea, high fever or purulent sputum occurs, a doctor or a pharmacist should be consulted immediately.

9. Interactions with other medicinal products and other forms of interaction

- None reported (2,5).

10. Fertility, pregnancy and lactation (2,5)

- Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.
- No fertility data available.

11. Effects on ability to drive and use machines (2,5)

- No studies on the effect on the ability to drive and use machines have been performed.

12. Undesirable effects

- If adverse reactions occur, a doctor or a pharmacist should be consulted.
- In sensitive persons, irritation of the gastric mucosa (5) and diarrhea (3) might occur.

13. Overdose

- No case of overdose has been reported (2,5).

14. Relevant biological activities

- Not required as per Egyptian guidelines for registration of herbal medicines.

15. Additional information

-

16. Date of compilation/last revision

13/11/2023

References

1.	www.powo.science.kew.org
2.	European Union herbal monograph on <i>Grindelia robusta</i> Nutt., <i>Grindelia squarrosa</i> (Pursh) Dunal, <i>Grindelia humilis</i> Hook. et Arn., <i>Grindelia camporum</i> Greene, herba (2012). EMA/HMPC/748220/2011. Committee on Herbal Medicinal Products (HMPC).
3.	PDR for Herbal Medicines (2002). Montvale, NJ: Medical Economics Company, 2 nd ed., ISBN 1-56363-361-2. 247-248
4.	Chevallier, A. (1996). The Encyclopedia of Medicinal Plants, Dorling Kindersley, London, UK.
5.	ESCOP Monographs (2015). Grindeliae herba - Grindelia . European Scientific Cooperative on Phytotherapy. Edited by Simon Mills and Roberta Hutchins. ISBN 978-1-901964-34-9.
6.	Spiteri, M. (2011). Herbal monographs including herbal medicinal products and food supplements. Department of Pharmacy, University of Malta. Set and printed by Print Right Ltd, Qormi.
7.	Timmermann, B., Wollenweber, E., Dörr, M., Armbruster, S., Valant-Vetschera, K. M. and Fuentes, E. R. (1994). External flavonoids in Two <i>Grindelia</i> species. <i>Z. Naturforsch.</i> , 49c , 395-397.
8.	Assessment report on <i>Grindelia robusta</i> Nutt., <i>Grindelia squarrosa</i> (Pursh) Dunal, <i>Grindelia humilis</i> Hook. et Arn., <i>Grindelia camporum</i> Greene, herba (2012). EMA/HMPC/748218/2011. Committee on Herbal Medicinal Products (HMPC).
9.	El-Shamy, A. M., El-Hawary, S. S., El-Shabrawy, A. O., El-Hefnawy, H. M. and Glasl, H. (2000). Essential oil composition of three <i>Grindelia</i> species. <i>Journal of Essential Oil Research</i> , 12 (5), 631-634. DOI: 10.1080/10412905.2000.9712175.
10.	Fraternale, D., Giamperi, L., Bucchini, A. and Ricci, D. (2007). Essential oil composition and antioxidant activity of aerial parts of <i>Grindelia robusta</i> from Central Italy. <i>Fitoterapia</i> , 78 (6), 443-5.
11.	Veres, K., Roza, O., Laczkó-Zöld, E. and Hohmann, J. (2014). Chemical composition of essential oils of <i>Grindelia squarrosa</i> and <i>G. hirsutula</i> . <i>Nat. Prod. Commun.</i> , 9 (4), 573-4. PMID: 24868887.
12.	Poudel, A., Dosoky, N. S., Satyal, P., Swor, K. and Setzer, W. N. (2023). Essential oil composition of <i>Grindelia squarrosa</i> from Southern Idaho. <i>Molecules</i> , 28 (9), 3854. DOI: 10.3390/molecules28093854.
13.	Krenn, L., Wollenweber, E., Steyrleuthner, K., Görick, C. and Melzig, M. F. (2009). Contribution of methylated exudate flavonoids to the anti-inflammatory activity of <i>Grindelia robusta</i> . <i>Fitoterapia</i> , 80 (5), 267-269.
14.	Nowak, S. and Rychlińska, I. (2012). Phenolic acids in the flowers and leaves of <i>Grindelia robusta</i> Nutt. and <i>Grindelia squarrosa</i> Dun. (Asteraceae). <i>Acta Pol. Pharm.</i> , 69 (4), 693-8. PMID: 22876612.